



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

intra-uterine growth aside from its function as a transfer system.

FREDERICK S. HAMMETT,

LYLE G. MCNEILE

COLLEGE OF PHYSICIANS AND SURGEONS,
UNIVERSITY OF SOUTHERN CALIFORNIA,
LOS ANGELES, CALIF.

THE EFFECT OF DRAINAGE ON SOIL ACIDITY

For the purpose of studying the effect of drainage on soil acidity, samples of soil were taken in October, 1916, from three of the experiment fields of the Purdue Agricultural Experiment Station. These fields are located near Westport, North Vernon and Worthington. The soils of these fields are all heavy silt loam, very low in organic matter and naturally poorly drained and quite acid in reaction. All of these fields have been thoroughly tile drained from three to five years. A portion of the Westport field is undrained and there are adjacent undrained, untreated areas alongside the North Vernon and the Worthington fields.

TABLE I

RELATIVE ACIDITY OF DRAINED AND UNDRAINED SOILS

Field and Soil Treatment	Lbs. CaCO ₃ Needed per 2,000,000 Lbs. Soil	
	Drained	Undrained
<i>Westport field:</i>		
Limestone.....	40 #	760 #
Limestone, phosphate and potash.....	30 #	360 #
Untreated.....	860 #	1,280 #
<i>North Vernon field:</i>		
Untreated.....	1,880 #	2,840 #
<i>Worthington field:</i>		
Untreated.....	740 #	1,600 #

Table I. shows the acidity of the soil as determined by the potassium nitrate method. Without entering into a discussion of the merits of different soil acidity methods, it may be said that on these soils, which are low in organic matter, there is no great difference in the degree of acidity shown by this method and the lime water and calcium salt methods. These results are consistent enough to indicate that drainage has a material influence on the acidity of soil of this type.

Farmers often refer to wet, poorly drained land as sour. While agricultural writers have placed little or no emphasis on such a correla-

tion, it is quite probable that soils in general will tend to become less acid when thoroughly drained, and vice versa; they will tend to become more acid when water-logged and poorly aerated. In testing soil acidity at different seasons of the year the results often vary quite a little in samples from the same plots of soil. These differences can not be attributed altogether to errors in sampling. The writer believes that at least part of the change of acidity is due to difference in aeration and moisture content of the soil at different seasons. Lipman and Waynick,¹ in an investigation of the effect of climate on soil properties, report that Maryland soil, which shows an acid reaction in its original location, when transported to Kansas or to California becomes neutral or slightly alkaline. It is quite probable that the better drainage and aeration of the soil when placed under less humid conditions could account very largely for the changes in reaction.

Considering SiO₂ an acid-forming oxide, practically all soils except those very high in the basic reacting elements, have a potentially great capacity for developing an acid reaction.

The writer believes that the constitution of the silicates of aluminum has more to do with injurious soil acidity than any other single factor. The acidity of aluminum silicates varies both with the relative proportion of SiO₂ to Al₂O₃ and with the amount of combined water in the silicate.² The weathering and changing of soil silicates under poorly drained or well-drained conditions would undoubtedly vary the constitution of the silicates and also vary the degree of soil acidity. It is quite true that certain types of well-drained sandy soils are acid. It is true also that a number of other factors besides drainage conditions affect soil acidity, but it is probable that the most acid soils are formed in poorly drained areas.

S. D. CONNER

INDIANA AGRICULTURAL EXPERIMENT STATION,
LAFAYETTE, IND.

¹ Lipman, C. B., and Waynick, D. D., *Soil Science*, Vol. I., No. 1, p. 5, 1916.

² Conner, S. D., "Acid Soils and the Effect of Acid Phosphate and Other Fertilizers upon Them," *Jour. Ind. and Eng. Chem.*, Vol. VII., No. 1, p. 35, 1916.